

Prevalence of Developmental Defects of Enamel in Children Visiting a University Hospital in Chennai - A Retrospective Study

Running title: Prevalence of developmental defects of enamel in children

KARTHIGA DEVI.G

*Saveetha Dental College and Hospitals,
Saveetha Institute of Medical and Technical Sciences,
Saveetha University Chennai- 600077.
Email ID: 151501066.sdc@saveetha.com*

GEO MANI

Reader

*Department of Pedodontics and Preventive Dentistry,
Saveetha Dental College and Hospitals,
Saveetha Institute of Medical and Technical Sciences,
Saveetha University
Chennai- 600077.
Email ID: geomani.sdc@saveetha.com*

PRADEEP D

Reader,

*Department of Oral and Maxillofacial Surgery,
Saveetha Dental College and Hospitals,
Saveetha Institute of Medical and Technical Sciences,
Saveetha University,
162,Poonamallee High Road, Chennai-600077,Tamilnadu, India.
Email id- pradeep@saveetha.com*

CORRESPONDING AUTHOR

GEO MANI

Reader

*Department of Pedodontics and Preventive Dentistry,
Saveetha Dental College and Hospitals,
Saveetha Institute of Medical and Technical Sciences,
Saveetha University, 162 Poonamallee High Road,
Chennai- 600077. Tamil Nadu, India.
Email ID: geomani.sdc@saveetha.com*

Article Info

Volume 83

Page Number: 2536 - 2551

Publication Issue:

July-August 2020

Article History

Article Received: 06 June 2020

Revised: 29 June 2020

Accepted: 14 July 2020

Publication: 25 July 2020

Abstract:

The developmental defects of the enamel (DDE) may be the alternation of enamel that may affect an area of one surface or may be widespread affecting all the surface of the teeth. Early diagnosis of developmental defects of the enamel can prevent the child from having an impact on oral health. The aim of this study was to determine the prevalence of developmental defects of enamel in children visiting a university hospital in the Chennai population. This was a retrospective study done under a University setting. It was seen that 334 patients had Developmental Defects of Enamel. The data was collected from June 2019 to March 2020. The data was tabulated and entered in excel and the data was analysed using SPSS package software. The overall prevalence of the DDE was found to be 47%. The most common deformity was Hypoplasia (24.3%) and the least common deformity was Molar Incisal Demineralisation and Amelogenesis Imperfecta (0.9%). The most common tooth affected was maxillary left central incisor (5.1%) and the least common deformity was found in mandibular right first premolar and primary dentition (0.3%). There was a statistically significant association between DDE and buccal surface of teeth. It indicates the need for educating the population about the risk factors for the DDE.

Keywords: Enamel defects, Hypoplasia, Diffuse Opacity, Demarcated Opacity, Molar Incisal Hypomineralisation, Children.

INTRODUCTION:

Oral health plays a very important role in the general well-being of individuals, and parents' behavior and attitudes influence the oral health of their children (Gurunathan and Shanmugaavel, 2016). The developmental defects of the enamel (DDE) may be defined as the alteration of enamel that may affect an area of one surface or may be widespread affecting all the surfaces throughout its full thickness (Subramanyam *et al.*, 2018). Opacity is a hypomineralization defect involving alteration in the translucency of enamel. Hypoplasia is a quantitative defect associated with a reduced thickness of enamel and appears as grooves or pits. These enamel defects can have a significant impact on esthetics, tooth sensitivity and occlusal function. They may be quantitative in nature that is manifested as a deficiency in adequate thickness of enamel or qualitative in nature as enamel opacities (Wright, 2000). Fluoride when present in optimal quantities is known to prevent caries by various mechanisms but more predominantly by

deposition of calcium fluoride crystals which is more resistant to demineralisation (Somasundaram *et al.*, 2015), ('Fluoride, Fluoridated Toothpaste Efficacy And Its Safety In Children - Review', 2018). DDE is associated with a wide spectrum of etiologic factors including systemic, genetic, local, and environmental conditions (Mihaela, Pasareanu and Maxim, 2011). Enamel defects have significant impact on oral health and esthetics and act as a predisposing factor for caries (Jeevanandan, 2017). Most epidemiological studies have shown that the frequencies of DDE are on rise in all populations streamlining their clinical significance and evidence for public health initiatives (Govindaraju and Jeevanandan, 2017a).

The prevalence of DDE ranged from 6.7% to 67.1% in the developed countries and from 27% to 66.2% in the developing countries. Chauhan *et al.*, (Chauhan and Chauhan, 2013) in 2013 assessed the prevalence and presentation of the DDE of healthy school children residing in hills of Himachal Pradesh, India, and reported to be 66.2%. Gisoo *et al.*, (Gisoo and Mohseni,

2010) (2010) found a significant association between systemic illness during the first 5 years of life and the prevalence of DDE. Mihaela et al., (Mihaela, Pasareanu and Maxim, 2011), reported a significant association between intake of drugs or chemicals during the first 5 years of life and prevalence of DDE. Enache et al., (Enache, Maxim and Păsăreanu, 2010) reported a significant association between nutritional status and prevalence of DDE.

The knowledge of the epidemiology of enamel defects is important in order to provide basic information within a community or country and between countries; and help in educating the population (Jeevanandan and Govindaraju, 2018). It is also important since it may contribute to the assessment and monitoring of environmental or systemic factors and for detecting possible etiological factors responsible for the occurrence of the enamel defects (Robles *et al.*, 2013).

Previously our team has conducted numerous cross sectional studies (Jeevanandan, 2017), (Govindaraju and Jeevanandan, 2017a), (Govindaraju and Jeevanandan, 2017b), (Somasundaram *et al.*, 2015), (Jeevanandan and Govindaraju, 2018), clinical trials (Govindaraju, Jeevanandan and Subramanian, 2017), (Ravikumar, Jeevanandan and Subramanian, 2017), (Panchal, Jeevanandan and Subramanian, 2019), (Christabel and Gurunathan, 2015), (Packiri, 2017), in-vitro studies (Gurunathan and Shanmugaavel, 2016), (Govindaraju and Gurunathan, 2017), case studies (Subramanyam *et al.*, 2018) and literature reviews ('Fluoride, Fluoridated Toothpaste Efficacy And Its Safety In Children - Review', 2018), (Nair *et al.*, 2018) over the past 5 years. Hence, this time, an attempt has been made to evaluate the prevalence of developmental defects of enamel in children visiting university in the Chennai population under 6-18 years old.

METHODS AND MATERIALS:

Ethical Approval:

The ethical approval for the retrospective study was obtained from the university ethical review board (Approval no: SDC/SIHEC/2020/DIASDATA/0619-0320).

Data analysis:

After reviewing the records of 86000 patients, a total of 334 records which satisfied the inclusion and exclusion criteria were included in the study. Simple random sampling was done to obtain the study sample. Population type was patients with age groups of 6 to 18 years. The patient records between June 2019 and March 2020 were reviewed and analysed. Cross verification was done by referring case sheets and photographs. All the incomplete & unreliable data was excluded. The Inclusion criteria was patients within age groups of 6 to 18 and patients with any systemic conditions or syndromes were excluded. The study was carried on in a university set up. The internal validity of the study was set by following proper sampling and also by strictly adhering to the protocols for examination of the case sheets for verification of cases with developmental enamel defects.

Statistical Analysis:

After Excel tabulation, the data was transferred to SPSS. The analysis was done using SPSS version 19. Descriptive statistics were used to correlate the data. The dependent variables were Hypoplasia, Diffuse opacity and Demarcated opacity. The independent variables were age and gender. The collected data was imported to SPSS and the chi square test was done. The level of significance was set at 0.05.

RESULTS AND DISCUSSION:

In the present study where 334 patients were examined, 50.30% were Males and 49.70% were Females [Figure 2].




Figure 2: Pie chart showing distribution of Gender of the patient. Black color represents the Male patients and Pink color represents the Female patients. Male (50.30%) and Females (49.70%) among the study population.

The overall prevalence of the DDE was found to be 47% (155 patients). The prevalence of DDE was found to be higher in Males (51.61%) when compared to Females (48.39%) [Figure 3].




Figure 3: Bar chart showing distribution of patients with Developmental Defects of Enamel (DDE), according to Gender of the patient. Black color represents the Male patients and Pink color represents the Female patients. The DDE was most common among the Male patients (51.61%) and least common among the Female patients (48.39%). Hence, the prevalence of DDE was higher among the Male patients compared to Female patients.

The most common tooth affected by the defect was common tooth was Mandibular Right First Premolar Maxillary Left Central Incisor (5.1%) and the least (0.3%) [Table 1].

Site (FDI System) that had DDE	FREQUENCY	PERCENTAGE (%)
0- No defect	179	53.6%
11/ 12/ 13	13/ 5/ 8	3.9%/ 1.5%
14/ 15	2/ 1	0.6%/ 0.3%
16	12	3.6%
21/ 22/ 23	17/ 2/ 8	5.1%/ 0.6%/ 2.4%
24/ 25	3 / 2	0.9%/ 0.6%
26	11	3.3%
31/ 32/ 33	4/ 3/ 3	1.2%/ 0.9%/ 0.9%
34/ 35	4/ 2	1.2%/ 0.6%
36	12	3.6%
41/ 42/ 43	5/ 2/ 2	1.5%/ 0.6%/ 0.6%
44/ 45	1/ 3	0.3%/ 0.9%
46	10	3.0%
52/ 53	2/ 1	0.6%/ 0.3%
54/ 55	3/ 3	0.9%/ 0.9%
62/ 63	1/ 2	0.3%/ 0.6%
73	2	0.6%
All Teeth	3	0.9%

TABLE 1: Table showing distribution of Developmental Defects of Enamel in individual teeth represented in FDI system of notation. Highest prevalence was seen in Maxillary left permanent central incisor.

The most common Site/Surface was the Labial surface (17.86%) and the least common affected surface was the mesial surface (0.3%) [Figure 4].




Figure 4: Pie chart showing distribution of patients with Developmental Defects of Enamel, according to surfaces affected by DDE of the patients. The most common Site/Surface was the Labial surface (17.66%) and the least common affected surface was the mesial surface (0.30%) . Hence, the prevalence of surfaces affected by the DDE was higher among the Labial surface compared to other surfaces of the teeth.

The most common deformity was hypoplasia (24.3%) Hypomineralisation and Amelogenesis Imperfecta and the least common deformity was Molar Incisal (0.9%) [Figure 5].




Figure 5: Pie chart showing distribution of patients with Developmental Defects of Enamel, according to the type of deformity in patients with DDE. The most common deformity was hypoplasia (24.3%) and the least common deformity was Molar Incisal Hypomineralisation and Amelogenesis Imperfecta (0.9%). Hence, the prevalence of type of deformity in patients with DDE was higher among hypoplasia compared to other types of deformity.

Hypoplasia (13.47%) was the highest type of deformity among the gender of the patients. Chi square test was performed and association between Type of deformity and Gender was not found to be statistically significant ($P=0.293$) [Figure 6].




Figure 6: Bar chart showing association between gender and various types of enamel deformity. X axis represents the number of patients with Type of Deformity. Y axis represents the number of patients with DDE. Hypoplasia was the highest type of deformity noticed among females (Pink) and Diffuse Hypoplasia was the highest type of deformity noticed among males (Black). Chi square test was performed and association between Type of deformity and Gender was not found to be statistically significant. P Value= 0.293 ($P>0.05$), hence not statistically significant.

association between DDE and Gender was found not to be statistically significant ($P = 0.655$) [Figure 7].

The DDE was most common among the Male patients (23.95%). Chi square test was performed and




Figure 7: Bar chart showing association between DDE and Gender. X axis represents the distribution of the number of patients with and without DDE . Y axis represents the number of patients with DDE. The DDE was most common among the Male patients (Black) compared to female patients (Pink). Chi square test was performed and association between DDE and Gender was found not to be statistically significant. P Value = 0.655 ($P>0.05$), hence not statistically significant.

Hypoplasia (11.96%) was the highest type of deformity among the buccal surface affected by the

DDE. Chi square test was performed and association between Surfaces and Type of deformity among patients with DDE was found to be statistically significant ($P = 0.000$) [Figure 8].




Figure 8: Bar chart showing association between Teeth surfaces and Type of deformity among patients with DDE. X axis denotes the Surfaces affected by DDE. Y axis denotes the type of deformity. Hypoplasia was the highest type of deformity among the buccal surface (Red) affected by the DDE. Chi square test was performed and association between Surfaces and Type of deformity among patients with DDE was found to be statistically significant. P value= 0.000 ($P > 0.05$), hence statistically significant.

The developmental defects of enamel can be defined as any alteration or disturbances during the process of

odontogenesis (Nelson *et al.*, 2010). According to their clinical appearance, DDE has been classified as Demarcated opacity, Diffuse opacity and Hypoplasia (Ravikumar, Jeevanandan and Subramanian, 2017). The absence of malnutrition and well-functioning healthcare during early childhood, may have contributed to a lower DDE prevalence. Frontal DDE led to aesthetic problems in every fourth case. Thus, the majority did not worry about the deviant appearance of their front teeth. Notably, the dentists were significantly more concerned than the patients. Consequently, the dentists must be careful not to transmit their opinion concerning appearance to the patients.

Isolated enamel opacity does not usually lead to an increased risk for caries; however it may be a significant esthetic problem. Enamel hypoplasia is not only a cosmetic problem, but it could also be a retentive area which favors bacterial plaque accumulation and results in increased caries susceptibility. Increased wear and tooth sensitivity lead to pain and discomfort for the child (Salanitri and Seow, 2013). Enamel hypoplasia may also be predictive of similar disturbances in the permanent dentition. Yellow - brown defects have lower Knoop hardness values and greater porosity than the white defects and normal enamel (Rodd, Boissonade and Day, 2007). Hypomineralized permanent first molars demonstrate changes in pulpal innervation and vascularity, and also show immune cell accumulation which is indicative of an inflammatory pulpal response (Jälevik and Klingberg, 2002). The

porous exposed subsurface enamel and the dentine may favor bacterial penetration into the dentine resulting in chronic inflammation of the pulp, which makes the use of local analgesia less efficient thus making the treatment painful, especially when immature teeth with large pulps are involved (Brogårdh-Roth and Matsson, 2011). Even intact enamel of hypomineralized molars may be very sensitive to air, cold or warmth and children often avoid brushing the sensitive molars (Jälevik and Klingberg, 2002). It is difficult for the children with intellectual disabilities to maintain good oral hygiene by themselves. They have impaired teeth - brushing ability and less patience due to their motor, sensor and learning disabilities (Slayton *et al.*, 2001). Children can be more susceptible to dental caries if they reside at home and their parents are comforting them with cariogenic snacks (Wong *et al.*, 2014). Oral health promotion programs should be aimed at centers and schools for children with intellectual disabilities and should include the regular use of oral health services, oral hygiene education and dietary counseling (Wong *et al.*, 2014).

In the present study, the prevalence of DDE in children aged 6-18 years was found to be 47%. On literature search, the least prevalence of DDE reported was 6.7% Pasareanu *et al.*, among 600 school children aged between 8 years and 11 years from Iasi (Păsăreanu and Florea, 2001) and the highest prevalence of DDE reported was 67.1% Sujat *et al.*, among 1,024 school children aged 16 years from the island of Penang, Malaysia (Sujak, Abdul Kadir and Dom, 2004).

In our study, the prevalence of defects was found to be higher in Males (51.61%) compared to Females (48.39%) which was similar to the findings of Hussian *et al.*, (Nik-Hussein *et al.*, 2018). In the study conducted by Ramesh *et al.*, the prevalence of defects was slightly higher in females (90.7%) as compared to males (87.9%) which was contradictory to our study (Ramesh *et al.*, 2011). The most common tooth affected by the

defect was Maxillary Left Central Incisor (5.1%) and the least common tooth was Mandibular Right First Premolar (0.3%) which was similar to the findings of Ravindran *et al.*, (Ravindran and Saji, 2016) and the report by Montero *et al.*, (Montero, Douglass and Mathieu, 2003) stated that the posterior teeth were twice most commonly affected which is contradictory to our present study.

The most affected surface was Labial surface (17.7%) and the least common affected surface was the mesial surface (0.3%) whereas Ruiz *et al.*, stated the defects were involving less than one-third of the tooth surface and incisal (33%) which is contraindicated to our study (Ruiz *et al.*, 2013).

Robes *et al.*, (Robles *et al.*, 2013) found that the most common type of DDE to be Demarcated opacity was not similar to the present study while our study states that the most common deformity was Hypoplasia (24.3%) and the least common deformity was Molar Incisal Hypomineralisation and Amelogenesis Imperfecta (0.9%).

The prevalence of Molar Incisal Hypomineralisation was found to be (0.9%) in the present study which was contraindicated to the study conducted by Gomez *et al.*, (Martínez Gómez *et al.*, 2012) reported that the 17.85% had MIH and Sonmez *et al.*, (Sönmez, Yıldırım and Bezgin, 2013) reported that the prevalence of MIH in their study was found to be 7.77%.

Vello *et al.*, (Velló *et al.*, 2010) and Faria *et al.* (Corrêa-Faria *et al.*, 2013) reported a significant association between the prevalence of the DDE and age which is not similar to our present study. Funakoshi *et al.*, (Funakoshi, Kushida and Hieda, 1981) reported the significant association between the tooth number, surface of the tooth with the DDE which is similar to the present study.

Although the clinical significance of enamel defects is well known, the pathogenesis of the defects is still being studied. While the range of environmental insults that can damage the enamel organ have been identified, the threshold for damage, and the relative susceptibility of the enamel organ at the various stages of development have not been well researched. . Therefore, despite major advancement in knowledge regarding the nature of the defects and the genes involved in enamel and dentine defects, further research is required. Additionally, as enamel defects are currently managed by treating the symptoms, future research should also focus on development of suitable techniques and aesthetic restorative materials that can bond effectively to defects enamel and dentine.

CONCLUSION:

In conclusion, the overall prevalence of the DDE was found to be 47%. The high prevalence of developmental enamel defects found in Male patients compared to Females, association between DDE, Type of deformity and Gender was not found to be statistically significant. association between Buccal Surfaces and Type of deformity among patients with DDE was found to be statistically significant. In view of the prevalence of DDE and the fact that enamel hypoplasia constitutes a risk factor for future ECC, enamel defects should be included as a dental health indicator in epidemiological studies of children.

REFERENCES:

1. Brogårdh-Roth, S. and Matsson, L. (2011) 'Molar-incisor hypomineralization and oral hygiene in 10-to-12-yr-old Swedish children born preterm', *European journal of oral*. Wiley Online Library. Available at: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1600-0722.2011.00792.x>.
2. Chauhan, D. and Chauhan, T. (2013) 'Prevalence of developmental defects of enamel in mixed and permanent dentition of 9 and 12 year old children of Himachal Pradesh, India: A cross sectional ...', *International Journal of Health & Allied*. ijhas.in. Available at: <http://www.ijhas.in/article.asp?issn=MPWK-0010;year=2013;volume=2;issue=3;spage=185;epage=188;aulast=Chauhan>.
3. Christabel, S. L. and Gurunathan, D. (2015) 'Prevalence of type of frenal attachment and morphology of frenum in children, Chennai, Tamil Nadu', *World J Dent*, 6(4), pp. 203–207.
4. Corrêa-Faria, P. *et al.* (2013) 'Developmental defects of enamel in primary teeth: prevalence and associated factors', *International journal of paediatric dentistry / the British Paedodontic Society [and] the International Association of Dentistry for Children*, 23(3), pp. 173–179.
5. Enache, R., Maxim, A. and Păsăreanu, M. (2010) 'Risk factors involved in the development of enamel defects', *J Roman Med Dents*, 14, pp. 71–74.
6. 'Fluoride, Fluoridated Toothpaste Efficacy And Its Safety In Children - Review' (2018) *International Journal of Pharmaceutical Research*, 10(04). doi: 10.31838/ijpr/2018.10.04.017.
7. Funakoshi, Y., Kushida, Y. and Hieda, T. (1981) 'Dental observations of low birth weight infants', *Pediatric dentistry*, 3(1), pp. 21–25.
8. Gisoo, F. F. and Mohseni, A. (2010) 'Prevalence study of etiologies of developmental defects of enamel of first permanent molar among six to seven years old children', *Curr Res Dent*. pdfs.semanticscholar.org. Available at:

<https://pdfs.semanticscholar.org/6a7b/42d46d9dea2ec9bf1367fb7428c886c83be9.pdf>.

9. Govindaraju, L. and Gurunathan, D. (2017) 'Effectiveness of Chewable Tooth Brush in Children-A Prospective Clinical Study', *Journal of clinical and diagnostic research: JCDR*, 11(3), pp. ZC31–ZC34.
10. Govindaraju, L. and Jeevanandan, G. (2017a) 'Comparison of quality of obturation and instrumentation time using hand files and two rotary file systems in primary molars: A single-blinded randomized ...', *European journal of. thieme-connect.com*. Available at: https://www.thieme-connect.com/products/ejournals/html/10.4103/ejd.ejd_345_16.
11. Govindaraju, L. and Jeevanandan, G. (2017b) 'Knowledge and practice of rotary instrumentation in primary teeth among indian dentists: A questionnaire survey', *Journal of International. jioh.org*. Available at: <http://www.jioh.org/article.asp?issn=0976-7428;year=2017;volume=9;issue=2;spage=45;epage=48;aulast=Govindaraju>.
12. Govindaraju, L., Jeevanandan, G. and Subramanian, E. (2017) 'Clinical Evaluation of Quality of Obturation and Instrumentation Time using Two Modified Rotary File Systems with Manual Instrumentation in Primary Teeth', *Journal of clinical and diagnostic research: JCDR*, 11(9), pp. ZC55–ZC58.
13. Gurunathan, D. and Shanmugaavel, A. K. (2016) 'Dental neglect among children in Chennai', *Journal of the Indian Society of Pedodontics and Preventive Dentistry*, 34(4), pp. 364–369.
14. Jälevik, B. and Klingberg, G. A. (2002) 'Dental treatment, dental fear and behaviour management problems in children with severe enamel hypomineralization of their permanent first molars', *International journal of paediatric dentistry / the British Paedodontic Society [and] the International Association of Dentistry for Children*, 12(1), pp. 24–32.
15. Jeevanandan, G. (2017) 'Kedo-S Paediatric Rotary Files for Root Canal Preparation in Primary Teeth - Case Report', *Journal of clinical and diagnostic research: JCDR*, 11(3), pp. ZR03–ZR05.
16. Jeevanandan, G. and Govindaraju, L. (2018) 'Clinical comparison of Kedo-S paediatric rotary files vs manual instrumentation for root canal preparation in primary molars: a double blinded randomised clinical trial', *European archives of paediatric dentistry: official journal of the European Academy of Paediatric Dentistry*, 19(4), pp. 273–278.
17. Martínez Gómez, T. P. *et al.* (2012) 'Prevalence of molar-incisor hypomineralisation observed using transillumination in a group of children from Barcelona (Spain)', *International journal of paediatric dentistry / the British Paedodontic Society [and] the International Association of Dentistry for Children*, 22(2), pp. 100–109.
18. Mihaela, M., Pasareanu, M. and Maxim, A. (2011) 'Etiological factors of enamel developmental defects of permanent teeth in children and adolescent', *Romanian J Oral Rehabil*, 3, pp. 72–78.
19. Montero, M. J., Douglass, J. M. and Mathieu, G. M. (2003) 'Prevalence of dental caries and enamel defects in Connecticut Head Start children', *Pediatric dentistry*, 25(3), pp. 235–

239.

20. Nair, M. *et al.* (2018) 'Comparative evaluation of post-operative pain after pulpectomy with k-files, kedo-s files and mtwo files in deciduous molars -a randomized clinical trial', *Brazilian Dental Science*, 21(4), pp. 411–417.
21. Nelson, S. *et al.* (2010) 'Dental caries and enamel defects in very low birth weight adolescents', *Caries research*, 44(6), pp. 509–518.
22. Nik-Hussein, N. N. *et al.* (2018) 'PREVALENCE OF DEVELOPMENTAL DEFECTS OF ENAMEL AMONG 16-YEAR-OLD CHILDREN IN MALAYSIA', *Annals of Dentistry University of Malaya*, 6(1), pp. 11–16.
23. Packiri, S. (2017) 'Management of Paediatric Oral Ranula: A Systematic Review', *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH*. doi: 10.7860/jcdr/2017/28498.10622.
24. Panchal, V., Jeevanandan, G. and Subramanian, E. (2019) 'Comparison of instrumentation time and obturation quality between hand K-file, H-files, and rotary Kedo-S in root canal treatment of primary teeth: A randomized controlled trial', *Journal of the Indian Society of Pedodontics and Preventive Dentistry*, 37(1), pp. 75–79.
25. Păsăreanu, M. and Florea, C. (2001) 'Risk factors involvement in enamel dental displasya', *J De Med Preven*. Citeseer. Available at: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.567.2245&rep=rep1&type=pdf>.
26. Ramesh, G. *et al.* (2011) 'Developmental defects of enamel in children of Davangere District and their relationship to fluoride levels in drinking water', *Asia-Pacific journal of public health / Asia-Pacific Academic Consortium for Public Health*, 23(3), pp. 341–348.
27. Ravikumar, D., Jeevanandan, G. and Subramanian, E. M. G. (2017) 'Evaluation of knowledge among general dentists in treatment of traumatic injuries in primary teeth: A cross-sectional questionnaire study', *European journal of dentistry*, 11(2), pp. 232–237.
28. Ravindran, R. and Saji, A. M. (2016) 'Prevalence of the developmental defects of the enamel in children aged 12-15 years in Kollam district', *Journal of International Society of Preventive & Community Dentistry*, 6(1), pp. 28–33.
29. Robles, M.-J. *et al.* (2013) 'Prevalence of enamel defects in primary and permanent teeth in a group of schoolchildren from Granada (Spain)', *Medicina oral, patologia oral y cirugia bucal*, 18(2), pp. e187–93.
30. Rodd, H. D., Boissonade, F. M. and Day, P. F. (2007) 'Pulpal status of hypomineralized permanent molars', *Pediatric dentistry*, 29(6), pp. 514–520.
31. Ruiz, L. A. *et al.* (2013) 'Prevalence of enamel defects in permanent teeth of patients with complete cleft lip and palate', *The Cleft palate-craniofacial journal: official publication of the American Cleft Palate-Craniofacial Association*, 50(4), pp. 394–399.
32. Salanitri, S. and Seow, W. K. (2013) 'Developmental enamel defects in the primary dentition: aetiology and clinical management',

Australian dental journal. Wiley Online Library, 58(2), pp. 133–40; quiz 266.

33. Slayton, R. L. *et al.* (2001) 'Prevalence of enamel hypoplasia and isolated opacities in the primary dentition', *Pediatric dentistry*, 23(1), pp. 32–36.
34. Somasundaram, S. *et al.* (2015) 'Fluoride Content of Bottled Drinking Water in Chennai, Tamilnadu', *Journal of clinical and diagnostic research: JCDR*, 9(10), pp. ZC32–4.
35. Sönmez, H., Yıldırım, G. and Bezgin, T. (2013) 'The prevalence and severity of molar incisor hypomineralization in a group of children living in Ankara Turkey', *Clin Dent Res*, 37(1), pp. 35–41.
36. Subramanyam, D. *et al.* (2018) 'Comparative evaluation of salivary malondialdehyde levels as a marker of lipid peroxidation in early childhood caries', *European journal of dentistry*, 12(1), pp. 67–70.
37. Velló, M. A. *et al.* (2010) 'Prenatal and neonatal risk factors for the development of enamel defects in low birth weight children', *Oral diseases*, 16(3), pp. 257–262.
38. Wong, H. M. *et al.* (2014) 'Risk Factors of Developmental Defects of Enamel-A Prospective Cohort Study', *PLoS ONE*, p. e109351. doi: 10.1371/journal.pone.0109351.
39. Wright, J. T. (2000) 'Normal formation and development defects of the human dentition', *Pediatric clinics of North America*, 47(5), pp. 975–1000.