

# Synthesis, Characterization and microbial studies of Cu (II), Co (II) and Ni (II) Complexes of Schiff base derived from 2-hydroxy Naphthaldehyde and Leucine

Aminu Ahmed<sup>1</sup>, Abdu M. Ali<sup>2</sup> and Kutama I. U<sup>3</sup>.

<sup>1</sup>Department of Chemistry, Kano University of Science and Technology, Wudil, Kano State, Nigeria

<sup>2</sup>Department of Chemistry, Jigawa State College of Education, Gumel, Nigeria

<sup>3</sup>Department of Chemistry, Kano University of Science and Technology, Wudil, Kano State, Nigeria  
aahmedtw64@yahoo.com, abdumohd92@gmail.com

## Article Info

Volume 82

Page Number: 13988 – 13993

Publication Issue:

January-February 2020

## Abstract

Schiff base ligand got from buildup of 2-hydroxy naphthaldehyde and leucine was incorporated and utilized for the arrangement of Cu (II), Co (II) and Ni (II) buildings. The blended ligand and buildings were dissected by decay temperature, solvency, attractive helplessness, molar conductance and infrared spectra. The disintegration temperatures of the edifices are in the scope of 200-247°C, molar conductance esteems are in the scope of 2.23-5.86 Ohm-cm<sup>2</sup> mol<sup>-1</sup>. Attractive defenselessness estimation demonstrated that all buildings are paramagnetic while dissolvability test uncovered that all edifices and ligand are solvent in DMSO. The expository information show the arrangement of 1:1 metal to ligand proportion, the readied ligand displayed tridentate coordination and shaped chelates of octahedral geometry. The ligand and metal chelates have been read for microbial action utilizing *Proteus vulgaris*, *Staphylococcus aureus* (microscopic organisms) and *Candida albicans* (growths) by circle dissemination technique. The outcomes demonstrated that metal buildings are increasingly dynamic contrasted and Schiff base ligand against the test life forms.

## Article History

Article Received: 18 May 2019

Revised: 14 July 2019

Accepted: 22 December 2019

Publication: 26 February 2020

**Keywords:** Characterization, Microbial studies, Schiff base, Synthesis

## I. INTRODUCTION

Schiff bases are intensify that comprise azomethine bunch (- HC=N-) and had been first announced by way of Hugo Schiff in 1864, [1].

The arrangements of Schiff base for the maximum component happen below corrosive or base catalysis or with warmth. Schiff base is crystalline sturdy, weakly (feebly) essential and some structure insoluble salts with solid acids. Schiff base generally co-ordinates by O particle of the deprotonated phenolic accumulating and the N molecule of azomethine gathering, [2].

The keeping potential of ligands is based upon the concept of iotas (which move approximately as

co-appointment destinations), steric element and electro antagonism. On the off threat that the ligand have beneficial accumulating, for instance, - OH or - COOH near the website online of buildup, a genuinely regular five or six membered chelate ring may be fashioned. Tridentate Schiff base ligands shape in addition solid buildings, [1], [3].

Change steel edifices of Schiff bases have pulled in a notable deal of enthusiasm because of their strong herbal physical games, as an example, antifungal, antibacterial and anticancer packages. Studies have verified that complexation of metals to Schiff base ligands improves the antimicrobial

and anticancer sporting activities of the ligands [4].

In this paper the metal buildings of Cu(II), Co(II) and Ni(II) with the Schiff base inferred from 2-hydroxynaphthaldehyde and leucine is depicted in this manner portraying the engineered utility and microbial sporting events of the base.

## II. MATERIALS AND METHODS

Every one of the reagents utilized are of explanatory evaluation and have been obtained from zayo sigma artificial restricted and were utilized legitimately moving forward without any more decontamination. Softening point and deterioration temperature were resolved making use of Gallenkamp dissolving factor contraption. Conductivity estimation become done making use of Jenway conductivity meter model 4010, even as appealing powerlessness estimations were finished utilising MBS MKI appealing weakness balance at 25oC.


### Readiness of Schiff base Ligand

The strategy announced by using [5] changed into applied. An solution of leucine(1.31g,zero.01mole) in 10mL delicate water containing sodium hydroxide(zero.04g) changed into made in 250mL round bottomed carafe. Ethanolic association of 2-hydroxy naphthaldehyde (1.72g, 0.01mole) turned into blanketed drop savvy with consistent blending and warmed underneath reflux for 5-hours.The response combo changed into cooled to room temperature, first-rate yellow inspire framed was separated off, washed with ethanol-water blend and dried in a desiccator over anhydrous calcium chloride.

### Arrangement of Schiff Base Metal (II) Complexes

Metal (II) chloride association (0.0009mole) (in 10mL water) was protected drop astute with regular mixing to an ethanolic association of the ligand (zero.5g, zero.0018mole) (in 10mL

ethanol), the combo become warmed underneath reflux for five-hours.The response blend was accepted to represent five-days at room temperature. The multiplied complex changed into separated off, washed with ethanol-water combination and dried in a desiccator over anhydrous calcium chloride [6].



**Scheme I:** Preparation of Metal (II) Complex

Assurance of Metal to Schiff base Ratio inside the edifices Using Job's Method of Continuous Variation "The amount of facilitated ligand within the metallic particle changed into resolved utilizing Job's method for continual range. 3-millimolar methanol preparations of the ligand and metal chloride have been readied. The accompanying ligand to steel salt (mL); 15:01, 13:03, eleven:05, 09:07, 07:09,05:eleven, 03:13, 01:15,00:sixteen had been taken from the ligand association and each one of the metal complex. An all out extent of 16ml was stored up (inside the above request) at some stage in the manner and mole a part of the ligand become determined in every mixture. The preparations of the steel chloride (clean) had been checked to were given wave length of most intense assimilation ( $\lambda_{max}$ ) for every metallic particle. The spectrophotometer changed into set at  $\lambda_{max}$  before taking the absorbance esteem [7]."

A plot of absorbance against mole fraction of the ligand and the number of coordinated ligand was determined using the following relation:

$$(1) \quad n = \frac{X_i}{1 - X_i}$$

Where:

$n$  = number of coordinated ligand at maximum absorbance

$X_i$  = mole fraction at maximum absorbance

### Results and Discussions

Table A: Physical properties of ligand and its metal (II) complexes

Compound	Colour	BM( $\mu_{\text{eff}}$ )	Melting point ( $^{\circ}\text{C}$ )	Decomposition Temperature ( $^{\circ}\text{C}$ )	Percentage Yield (%)
Ligand	Bright Yellow	—	134	—	75
[Co L(H <sub>2</sub> O) <sub>3</sub> ]	Brown	4.42		200	45
[Cu L(H <sub>2</sub> O) <sub>3</sub> ]	Green	1.62		247	54
[Ni L(H <sub>2</sub> O) <sub>3</sub> ]	Yellowish Green	2.60		233	62

Table B: Solubility tests of ligand and its metal (II) complexes

Compound	Distilled Water	Me OH	Et O H	Chloroform	Ace tone	n-He xane	D MS O
Ligand	IS	S	S	IS	S	IS	S
[Co L(H <sub>2</sub> O) <sub>3</sub> ]	IS	S	S	IS	S	IS	S

[Cu L(H <sub>2</sub> O) <sub>3</sub> ]	IS	SS	SS	IS	SS	IS	S
[Ni L(H <sub>2</sub> O) <sub>3</sub> ]	IS	SS	SS	IS	SS	IS	S

KEY: S = Soluble SS = Slightly Soluble IS = Insoluble

Table C: Conductivity measurement of complexes in  $3 \times 10^{-3}$  DMSO

Complex	Concentration(mol/dm <sup>3</sup> )	Specific Conductance (Ohm <sup>-1</sup> cm <sup>-1</sup> )	Molar Conductance (Ohm <sup>-1</sup> cm <sup>2</sup> mol <sup>-1</sup> )
[Co L(H <sub>2</sub> O) <sub>3</sub> ]	$3 \times 10^{-3}$	$10.46 \times 10^{-6}$	3.49
[Cu L(H <sub>2</sub> O) <sub>3</sub> ]	$3 \times 10^{-3}$	$6.69 \times 10^{-6}$	2.23
[Ni L(H <sub>2</sub> O) <sub>3</sub> ]	$3 \times 10^{-3}$	$17.58 \times 10^{-6}$	5.86

Table D: Infrared spectral data of the ligand and its metal (II) complexes

Compound	$\nu$ (OH)/H <sub>2</sub> O cm <sup>-1</sup>	$\nu$ (C=N) cm <sup>-1</sup>	$\nu$ (M-N) cm <sup>-1</sup>	$\nu$ (M-O) cm <sup>-1</sup>
Ligand	3056	1617	-	-
[Co L(H <sub>2</sub> O) <sub>3</sub> ]	3257	1621	672	466
[Cu L(H <sub>2</sub> O) <sub>3</sub> ]	3052	1647	561	442
[Ni L(H <sub>2</sub> O) <sub>3</sub> ]	3164	1625	633	432

Table E: Antibacterial and antifungal activities of ligand and its complexes

Compound	Inhibition zone(mm)								
	<i>Proteus vulgaris</i>			<i>Staphylococcus aureus</i>			<i>Candida albican</i>		
	2	1		1	1		1	1	
Concentration(mg/mL)	0	0	5	20	0	5		0	5
	0	0	0	0	0	0	200	0	0
Ligand	1		N						N
	0	8	I	12	8	ZI	11	9	I
[Cu L(H <sub>2</sub> O) <sub>3</sub> ]	1	1			N				N
	7	2	8	9	I	ZI	14	7	I
[Co L(H <sub>2</sub> O) <sub>3</sub> ]	1	1							
	6	0	7	13	8	6	15	9	7
[Ni L(H <sub>2</sub> O) <sub>3</sub> ]			N					N	N
	1		Z			N	NZ	Z	Z
	3	9	I	12	8	ZI	I	I	I
Control	Gentamycin e 15			Polymyxin 12			Ketoconazole 25		

### III. Discussions

The Schiff base ligand turned into installation through buildup of leucine and a pair of-hydroxy naphthaldehyde to get amazing yellow valuable stone with high return (seventy five%) and softening reason of 134oC (desk A).The metal edifices were set up by the reaction of steel (II) chloride with the Schiff base ligand and are of different hues, their fee yields are inside the scope of forty five-sixty two%.

The decay temperature of the metallic (II) buildings is better than that of liquefying reason of the ligand showing that complexation has occurred [6].

Attractive helplessness estimation at 25oC exposed that the metallic (II) buildings are

paramagnetic in nature. Likewise, the powerful attractive minute features are interior anticipated scope of octahedral edifices; this is predictable with hypothetical desire and what turned into accounted for within the writing [8]. The attractive snapshot of Co (II) complex is four.42BM inside the scope of four.3-5.0BM, Ni (II) complicated is two.6BM inside the scope of two.Five-3.5BM and Cu (II) complex is 1.Sixty two BM (1.7-2.2BM) which is the attribute of 3d9 [8] , [9].

The aftereffects of dissolvability check (desk B) of the Schiff base and edifices in diverse dissolvable exposed that every one the homes and Schiff base ligand are solvent in DMSO, insoluble in subtle water, chloroform and n-hexane, particularly dissolvable in methanol, ethanol and CH<sub>3</sub>)<sub>2</sub>CO with exemption of ligand and Co(II) complicated which are dissolvable.

All the edifices have molar conductance (desk C) inside the scope of two.23-five.86 ohm-1 cm<sup>2</sup>mol-1 showing that the homes are non-electrolytes [9] , [10].

The infrared phantom results (desk D) of the ligand suggests a band at1617 cm-1 allotted to azomethine ν (C=N) vibration, this affirms buildup between amino collecting of leucine and aldehyde accumulating of 2-hydroxy naphthaldehyde within the improvement of Schiff base [5] .The medium force pinnacle of ingestion in the Schiff base ligand became seen at 3056 cm-1, because of naphthaldehyde hydroxyl bunch ν (OH) [11]. This pinnacle of assimilation vanishes within the IR spectra of the buildings which display coordination thru oxygen iota of the hydroxyl gathering of naphthaldehyde. The band in the scope of 1621-1647 cm-1 was likewise seen in all of the metal (II) buildings which are inferable from the coordination of nitrogen iota of azomethine gathering to the metallic particles. New band confirmed up in the spectra of the buildings inside the scope of 561-672 cm-1 and 432-466 cm-1 which reveal ν (M - N) and ν (M - O) vibrations personally asserting coordination of

Schiff base ligand to the metal debris thru nitrogen and oxygen atoms.

The metal to Schiff base share become resolved utilizing Job's approach for consistent range which endorsed 1:1 ligand to metal share in all the homes dissected.

The antibacterial action (table E) of the ligand and metallic edifices turned into completed on two microorganisms segregates: *Proteus vulgaris* and *Staphylococcus aureus* utilizing circle dispersion method. The distance throughout of dilemma sector (mm) was expected for each intensify, the steel edifices have been visible as steadily dynamic contrasted with Schiff base ligand towards the take a look at creatures under indistinguishable exploratory situations. The enlargement in action of the steel chelates is because of the effect of the metallic particle in standard cellular procedure as make clear with the aid of Tweedy's hypothesis. Chelation decreases the extremity of the metallic particle due to the quilt of ligand orbital and fractional sharing of fantastic fee of the steel particle with the giver gathering. It moreover increments  $\pi$ -electron delocalization in the whole chelate jewelry this is formed in the course of coordination. Consequently improves lipophilic person of the focal steel iota and builds the hydrophobic character and lipo-solvency of the complex assisting its infiltration through the lipid layers of the mobile movie in this manner deactivating breath method inside the small scale life forms [6]. The antifungal movement (desk E) of the Schiff base and its metallic buildings have been examined on one parasitic seclude, *Candida albican* utilising circle dispersion method. The outcomes suggest that metallic homes have been greater dynamic than the Schiff base ligand with exemption of nickel complicated that is dormant towards the check life shape. The antifungal motion was additionally clarified via Tweedy's chelation speculation [ 6]. Some edifices and ligand display no restraint area at low cognizance

(50mg/l) in opposition to chose microbes and growths. The inhibitory depth of the homes turned into contrasted and that of the anti-infection retailers (*Gentamycine*, *Polymyxin* and *Ketoconazole*). The buildings display feasible motion towards the test dwelling beings; subsequently, they can be applied as option in evaluation to anti-infection marketers

#### IV. Conclusion

The metal (II) complexes were synthesized and analyzed by solubility, melting point/decomposition temperature, magnetic susceptibility molar conductance, infrared spectra. The metal to ligand ratio is 1:1 with the formula [ML], where M represents Cu(II),Co(II) and Ni(II) ions and L represents the Schiff base ligand.

#### References

- [1] R.R.Gupta, Kumar &V. Gupta, "Heterocyclic Chemistry I," pp.230-234, 1998.
- [2] B.K.Al-Salami, M.H.Mohammed, &okay.A. Askar, "mix and Characterization of latest Schiff Bases Derived From 2-hydroxybenzaldehyde and Amino Acids and Their Vanadyl Complexes," study Journal of Pharmaceutical, organic and Chemical Sciences Vol.5, pp.1457-1474, 2014.
- [3] B.Okay.Panda, &A.Chakravorty, "Spectroscopic places of inorganic and organometallic blends," Journal of Organometallic Chemistry Vol.690, pp.3169-3175, 2005.
- [4] M.Nasser, N.M.Hosny, &F.I.Dossoki, "Schiff Base Complexes Derived from 2-Acetylpyridine, Leucine and a few metallic Chlorides: Their education, Characterization and bodily residences," J. Chem. Eng. Vol.Fifty three, pp.2567-2572, 2008.
- [5] L.Lekha, k. Kanmaniraja, G.Rajagopal, D.Sivakumar and D.Easwaramoorthi, "organization, apparition depiction and antimicrobial assessment of Schiff Base ligand acquired from amino hazardous and its encouraging steel houses," worldwide Journal of Chemical and Pharmaceutical Sciences Vol. 4, pp.Forty 8-fifty four, 2013.

- [6] D.A. Skoog, D.M. West & F.J. Holler, "Nuts and bolts of Analytical Chemistry, "5th Ed. Saunders: new york, pp.525-528, 1988.
- [7] okay.Mounika, B.Anupama, J.Pragathi, & C.Gyanakumari, "Synthesis, Characterization and organic exercise of a Schiff Base Derived from three-Ethoxy Salicylaldehyde and a few Amino Benzoic unsafe and its Transition metal Complexes," J. Of Sci. Res. Vol.2, pp.513-524, 2010.
- [8] H.A.Nora, "Synthesis Characterisation and organic recreation of Cu(II), Co(II), Mn(II), Fe(II) UO<sub>2</sub>(VI) Complexes with a new Schiff Base Hydrazone: O-Hydronyacepophenone-7-Chloro-four-Quinoline Hydrazone" Open access Molecular Journal Vol.Sixteen, pp.8629-8645, 2011.
- [9] A.M.Shaker, L.A.E.Nassr, M.M.S.Adam, and that i.M.A.Mohamed, "combine, Characterization and Spectrophotometric experiences of Seven Novel Antibacterial Hydrophilic Iron(II) Schiff Base Amino Acid Complexes," Journal of the Korean Chemical Society Vol.Fifty seven, pp.560-568, 2013.
- [10] I.Grace, , E.Iorkpiligh, and T. Isaac, "combine, Characterization and Antimicrobial stories of Mn(II) Co(II) AND Zn(II) Schiff Base Complexes Derived from L-Arginine and 2-Hydroxy-1-Naphthaldehyde," global Journal of Science and study Vol.4, pp.979-982, 2015.